



Original Reports

DACC Resting State Functional Connectivity as a Predictor of Pain Symptoms Following Motor Vehicle Crash: A Preliminary Investigation

Jacklynn M. Fitzgerald,^{*} Emily L. Belleau,^{†,‡} Lauren E. Ehret,[§] Colleen Trevino,[¶] Karen J. Brasel,^{||} Christine Larson,^{**} and Terri deRoon-Cassini^{¶,1}

^{*}Department of Psychology, Marquette University, Milwaukee, Wisconsin, [†]Department of Psychiatry, McLean Hospital, Belmont, Massachusetts, [‡]Harvard Medical School, Boston, Massachusetts, [§]Rogers Behavioral Health, Minneapolis, Minnesota, [¶]Division of Trauma & Critical Care, Department of Surgery, Medical College of Wisconsin, Milwaukee, Wisconsin, ^{||}Oregon Health & Science University, Portland, Oregon, ^{**}Department of Psychology, University of Wisconsin-Milwaukee, Milwaukee, Wisconsin

Abstract: There is significant heterogeneity in pain outcomes following motor vehicle crashes (MVCs), such that a sizeable portion of individuals develop symptoms of chronic pain months after injury while others recover. Despite variable outcomes, the pathogenesis of chronic pain is currently unclear. Previous neuroimaging work implicates the dorsal anterior cingulate cortex (dACC) in adaptive control of pain, while prior resting state functional magnetic resonance imaging studies find increased functional connectivity (FC) between the dACC and regions involved in pain processing in those with chronic pain. Hyper-connectivity of the dACC to regions that mediate pain response may therefore relate to pain severity. The present study completed rsfMRI scans on N = 22 survivors of MVCs collected within 2 weeks of the incident to test whole-brain dACC-FC as a predictor of pain severity 6 months later. At 2 weeks, pain symptoms were predicted by positive connectivity between the dACC and the premotor cortex. Controlling for pain symptoms at 2 weeks, pain symptoms at 6 months were predicted by negative connectivity between the dACC and the precuneus. Previous research implicates the precuneus in the individual subjective awareness of pain. Given a relatively small sample size, approximately half of which did not experience chronic pain at 6 months, findings warrant replication. Nevertheless, this study provides preliminary evidence of enhanced dACC connectivity with motor regions and decreased connectivity with pain processing regions as immediate and prospective predictors of pain following MVC.

Perspective: This article presents evidence of distinct neural vulnerabilities that predict chronic pain in MVC survivors based on whole-brain connectivity with the dorsal anterior cingulate cortex.

© 2020 by United States Association for the Study of Pain, Inc.

Key words: Motor vehicle accident, traumatic injury, pain, functional magnetic resonance imaging, resting state, cingulate cortex.

Received November 12, 2018; Revised March 26, 2020; Accepted July 25, 2020.

¹These authors shared senior authorship.

Disclosures: All authors report no biomedical financial interests or potential conflicts of interest.

Funding: This work was supported by funds obtained through the Medical College of Wisconsin Injury Research Center funded by the CDC (R49/CE001175).

Address reprint requests to Jacklynn Fitzgerald, PhD, Department of Psychology, Marquette University, Cramer Hall, 328N, PO Box 1881, Milwaukee, WI 53201-1881 E-mail: jacklynn.fitzgerald@marquette.edu 1526-5900/\$36.00

© 2020 by United States Association for the Study of Pain, Inc.

<https://doi.org/10.1016/j.jpain.2020.07.002>

Motor vehicle crashes (MVCs) are a leading cause of traumatic injuries in the United States—second only to falls—and account for over 20% of all severe injuries that require hospital care.²¹ While a host of negative outcomes accompany MVCs, chief among these is the experience of severe pain symptoms, which affect up to 80% of MVC survivors immediately after injury.⁹ In terms of the long-term prognosis of pain outcomes, there is considerable variability. Large-scale and population-based studies demonstrate that anywhere from 12 to 40% of MVC

2 The Journal of Pain

survivors continue to suffer from pain months after injury.^{47,57} As pain symptoms become chronic, defined as pain that is maintained at least 6 months after injury, risk for diminished quality of life and psychiatric illness increases.^{3,24} Therefore, deciphering who is at risk for the emergence of chronic pain after injury provides an opportunity to intervene and positively affect overall health of MVC survivors. Currently, however, relatively little is known about what factors moderate the relationship between MVC-related injury and the development of chronic pain, demonstrating a need for more research to precisely elucidate which factors predict heterogeneity of pain outcomes in this population.¹⁶

Neurobiologically, the acute experience of pain involves activation in numerous discrete brain regions.⁶² During injury, pain signals are sent from the periphery to the brain by way of the spinal cord and brainstem, which subsequently transmit these signals to the thalamus.⁶ This information is then sent to the somatosensory cortex, involved in deciphering the location and intensity of incoming pain signals.^{2,13,59} The amygdala and insula, which are involved in salience detection, processing, and experience of emotion,^{38,53,58} are also active and contribute to overall pain processing.^{33,73}

In addition to these regions, which are principally involved in the appraisal of pain as a salient event,⁵² the spinothalamic system directly innervates higher cortical regions to manage and control the pain response, with a chief target being the dorsal anterior cingulate cortex (dACC).^{44,72} The dACC receives direct^{1,25} and indirect^{10,49,74} connections from the brainstem, in addition to possessing a bilateral connection with the amygdala.^{34,61,68,81} Based on this organization and in the context of acute pain response, the dACC has been identified as a critical region involved in integrating incoming signals governing the perception of pain (eg, originating from the periphery by way of the brainstem) with those involved in salience detection and the generation of negative affect (eg, originating from the amygdala).^{54,72,77} Given the dACC's broader role in response inhibition and action planning,^{12,27} the principal role of the dACC in response to pain is believed to center on integrating pain and affective signals in order to mediate adaptive control of this experience.⁷² This is supported by evidence from primate studies showing that the dACC is active in response to pain as well as when animals are fleeing from pain.^{41,43,50} This suggests a role for the dACC in mediating the motivational response to pain. Humans with lesions in the cingulate cortex report reduced affective responses to pain without altering their perception of incoming pain signals, further suggesting that the dACC is involved in affective response to pain.³² Evidence from functional magnetic resonance imaging (fMRI) studies in healthy volunteers also shows that perceived controllability of feeling pain tracks linearly with dACC engagement when pain is administered.⁷⁰ In the context of pain outcomes following MVCs, engagement of the dACC may therefore influence long-term pain prognosis given its role in pain regulation.

Indeed, there is evidence of altered dACC involvement in chronic pain based on data from cross-sectional fMRI studies examining dACC activation in individuals

DACC Functional Connectivity and Pain

with fibromyalgia and chronic low back pain. Individuals with these conditions exhibit greater activation of the dACC during nociceptive processing.^{19,35,66} Augmented engagement of the dACC during pain processing may compensate for greater anticipation and attention toward pain,¹⁹ as evidenced by the fact that dACC activation positively relates to greater subjective ratings of pain in these samples⁶⁶ and symptoms of pain catastrophizing—or characterizing pain as awful, horrible and unbearable.³⁷ As pain processing involves many brain regions, functional connectivity (FC) studies have been useful for understanding alterations within larger networks, with specific focus on altered spontaneous low-frequency (<0.1 Hz) fluctuations between regions at rest. This work also demonstrates altered connectivity between the ACC and pain processing regions in those with chronic pain. For instance, Cifre et al found greater FC between the ACC and insula, and reduced ACC-amygdala and ACC-brainstem FC in those with fibromyalgia compared to healthy controls.¹⁷ Enhanced ACC-insula connectivity may reflect greater interoceptive awareness of pain, while decreased ACC-amygdala and ACC-brainstem connectivity may represent abnormal bottom-up signaling of pain sensations and/or reduced descending modulation of pain.¹⁷ Conversely, decreased FC between the insula and dACC is associated with greater reduction of pain symptoms when patients with fibromyalgia were administered milnacipran, a selective serotonin and norepinephrine reuptake inhibitor for their pain symptoms.⁷¹ Together, this work demonstrates that greater FC between the dACC and pain processing regions (eg, insula, thalamus, brainstem) characterizes chronic pain response, but that this FC is reversed when pain symptoms are treated. Based on this work, functional neurocircuitry between the dACC and brain regions involved in acute pain processing may be related to long-term pain outcomes; however, we are unaware of a study that has examined dACC-FC as a predictor of prospective pain severity following MVCs.

The current study tested whole-brain dACC-FC as a predictor of chronic pain symptoms assessed 6 months after a MVC. Pain ratings and rsfMRI were collected in-person acutely (eg, within 2 weeks of the MVC) and pain ratings were collected again in-person 6 months later at a follow-up visit. We hypothesized that pain ratings across the 2 timepoints would be positively related to one another. In addition and based on prior neuroimaging work, we hypothesized that baseline measures of rsfMRI, specifically positive connectivity of the dACC with regions involved in acute pain processing (eg, the brainstem, thalamus, somatosensory cortex, amygdala, and insula) would predict greater pain symptoms at baseline and prospectively predict greater pain symptoms at 6 months.

Methods

Participants and Procedures

Participants were recruited from the emergency department of the Level 1 trauma center at Froedtert

Hospital/Medical College of Wisconsin in Milwaukee, Wisconsin. Participants were not admitted to long-term hospital care but were discharged following completion of acute care within the emergency department (ED). Inclusion criteria included: 1) MVC as mechanism of injury, 2) between the ages of 18 and 65, 3) ability to read and write English, and 4) ability to give informed consent. Participants were excluded if they had a head injury that resulted in loss of consciousness (Glasgow Coma Scale score <13 on ED arrival). Additional exclusion criteria specific to fMRI scanning included: 1) presence of ferromagnetic material within the body, 2) pregnancy or actively trying to become pregnant, 3) fear of enclosed spaces (eg, claustrophobia), and 4) inability to lie still for up to 1 hour. Participants were recruited over the phone from an ED discharge census and provided consent for participation when they arrived for rsfMRI testing and self-report scales 2 weeks after injury at a baseline appointment; self-report scales were again collected in-person 6 months later at follow-up. The Institutional Review Board at the Medical College of Wisconsin approved all study procedures and participants were monetarily compensated for their time.

Self-Report Measures and Analysis

Participants completed a self-report index of pain at both baseline and 6-month visits using the Visual Analog Scale for Pain (VAS Pain),⁵⁶ a reliable and valid measurement for the reporting of unidimensional pain^{22,29} and that has been used as a primary outcome measure of recurrent pain.¹⁸ Upon presentation of the VAS Pain, participants were asked to rate how much pain they were currently experiencing on a continuous scale of 0 to 10 (0 = no pain, 10 = worst pain). In addition to the VAS Pain, all participants also completed a self-report rating of post-traumatic stress symptoms at both visits using the Impact of Events Scale-Revised (IES-R),⁷⁸ given high comorbidity between pain symptoms and post-traumatic stress disorder (PTSD).⁷ The IES-R asks participants to rate how distressed they feel by common posttraumatic stress symptoms. In addition to clinical measures, participants were also asked about medications they were taken at the time of the fMRI assessment to determine opioid use for pain management

rsfMRI Acquisition, Preprocessing, and Analysis

All participants completed a 6-minute resting state scan during fMRI (eg, rsfMRI) at their baseline appointment approximately 2 weeks after injury. During the scan, participants viewed a white crosshair displayed on a black background and were instructed to keep their eyes open. Scanning was performed on a 3.0 Tesla short bore GE Signa Excite MRI system at the Medical College of Wisconsin. Functional T2*-weighted echoplanar images were collected in a sagittal orientation with the following parameters: repetition time (TR)/echo time (TE) = 2,000/25 ms; FOV = 24 mm; matrix = 64 × 64; flip angle = 77°; slice thickness = 3.5 mm. A high-resolution T1-weighted anatomical image was also acquired for co-registration with the following parameters: TR/TE = 8.2/3.2 ms; FOV = 240 mm; matrix = 256 × 224; flip angle = 12°; voxel size = 0.9375 × 0.9375 × 1 mm.

Individual functional images were analyzed using the CONN functional connectivity toolbox⁷⁹ and images were preprocessed according to standard procedures. Briefly, images underwent spatial realignment, slice-time correction, structural segmentation and normalization, and motion correction. As small head movements can cause spurious noise and distance-dependent changes in signal correlations,^{64,65} frame-wise displacement (FD) was computed to rule out confounding effects of motion. Volumes with FD >0.2 mm (plus 1-back and 2-forward neighboring volumes) were “scrubbed” (eg, removed from analysis) and subjects with >3 mm or 3 degrees of rotational cumulative movement were dropped from analysis. Images were normalized to the Montreal Neurological Institute (MNI) template and smoothed with a 4 mm³ Gaussian kernel. To isolate rsfMRI signal, resulting data were bandpass filtered at 0.01 to 0.10 Hz, while signal from cerebrospinal fluid and white matter along with motion realignment parameters were entered as regressors of no-interest to control for these effects during scanning.

For whole-brain dACC-seeded analyses, an anatomical dACC mask was created from the ACC mask in the AAL atlas.^{55,76} This ACC mask was edited using fslview (FSL v.5.0.9⁴⁵) to exclude rostral ACC, such that ACC at and below the genu of the corpus callosum was excluded in-line with dACC versus rostral ACC boundaries.¹² Fig 1 illustrates spatial location of the dACC mask used as the seed region.




Figure 1. Dorsal anterior cingulate cortex mask.

4 The Journal of Pain

We first examined the relationship between dACC whole-brain FC and pain symptoms at baseline, controlling for age, gender, opioid administration at baseline (dichotomous variable), and time since injury for the baseline appointment. Second, we examined the relationship between dACC whole-brain FC and pain symptoms at 6 months, controlling for these same covariates in addition to pain symptoms at baseline and PTSD symptom severity at 6 months (eg, IES-R) given high comorbidity between the development of chronic pain and PTSD.^{23,60,75} Age and gender covariates were included given wide range in age distribution and unequal gender distribution. Significant effects were examined using a height threshold of $P < .001$ (uncorrected) and cluster threshold of $P < .05$ corrected for multiple comparisons across the entire brain based on a false discovery rate in-line with recent recommendations^{26,63,80} and identical to other published methods.^{20,40,48} To illustrate effects, connectivity values were extracted from significant clusters and input into SPSS (Version 25.0).

Results

Participants

Based on inclusion/exclusion criteria, a total of 22 participants consented to and were included in the study. Participants were between the ages of 18 and 62 ($M = 31.82$, $SD = 11.28$); $n = 17$ participants were female (77%). In terms of race and ethnicity, $n = 11$ (50%) were Caucasian, $n = 8$ (36%) African-American, $n = 2$ (9%) Hispanic, and $n = 1$ (5%) unknown. As a result of the MVC, 12/22 (54.55%) experienced injuries involving more than one area on the body, while the location of injuries included the back (12/22; 54.55%), neck (9/22; 40.91%), head (4/22; 18.18%), arms (3/22; 13.64%), hips (2/22; 9.09%), legs (2/22; 9.09%), chest (1/22; 4.55%), and abdomen (1/22; 4.55%). In terms of concomitant conditions, at the time of admittance into the ED, 16/22 (72.73%) had no current medical conditions, 3/22 (13.64%) had asthma, 1/22 (4.55%) had high cholesterol, 1/22 (4.55%) had hypertension, and 1/22 (4.55%) had anemia. No participants had a history of chronic pain prior to the injury.

At baseline, participants were asked to rate the severity of their pain associated with the injuries sustained during the MVC. Pain ratings ranged from 0 to 5.50 ($M = 2.57$, $SD = 1.83$) at baseline, and 0 to 7.00 ($M = 1.34$, $SD = 1.80$) at 6 months. The number of participants who reported pain 6 months after injury was 12/22 (54.55%) as qualified by a pain rating > 0 , indicating presence of pain. A total of 7/22 (31.82%) participants reported either identical pain severity compared to baseline or an increase in pain severity at 6 months. Pain ratings at baseline were positively correlated to pain ratings at 6 months ($r(20) = 0.44$, $P = .04$). A total of 6 participants (27%) were taking opioids at baseline for pain management.

As time since injury may affect pain severity ratings, we also assessed the correlation between time since

DACC Functional Connectivity and Pain

injury for both baseline and 6-month appointments and pain severity ratings. Two time since injury measures were calculated using difference scores reflecting: 1) baseline data collection date—date of injury, and 2) 6-month data collection date—date of injury. Time since injury for baseline assessments averaged 13.64 ± 2.87 days, and time since injury for the 6-month assessments averaged 202.30 ± 17.97 days. There was a trending relationship between time since injury at the baseline assessment and pain severity ratings at that time ($r(20) = 0.41$, $P = 0.06$), although time since injury for 6-month assessments and pain severity ratings at 6 months were not associated with one another ($r(20) = -0.04$, $P = .87$).

Symptoms of PTSD ranged from 0.58 to 8.08 ($M = 4.47$, $SD = 2.15$) at baseline to 0 to 9.75 ($M = 2.50$, $SD = 2.61$) at 6 months. Severity of pain and PTSD symptoms were not related at either baseline ($r(20) = 0.15$, $P = .52$) or 6 months ($r(20) = 0.39$, $P = .07$).

rsfMRI Predictors of Pain at Baseline

Pain symptoms at baseline were predicted by positive connectivity between the dACC and a cluster traversing the precentral gyrus/primary motor cortex (M1) and premotor cortex (peak MNI: 30, -14, 70; $Z = 7.45$; volume = 1,440 mm³; $P = .004$ false discovery rate-corrected). Spatial location of significant FC cluster in the precentral gyrus is displayed in Fig 2A.

rsfMRI Predictors of Pain at 6 Months

Pain symptoms at 6 months were predicted by negative connectivity between the dACC and the precuneus (peak MNI: -6, -74, 32; $Z = 5.25$; volume = 1,272 mm³; $P = .004$ corrected). Spatial location of significant FC cluster in the precuneus is displayed in Fig 2B.

Discussion

The present study tested resting state FC of the dACC as a predictor of pain severity 6 months following a MVC. Several important findings emerged from this investigation: first, pain severity at baseline was predicted by positive connectivity between the dACC and precentral gyrus, while pain severity at 6 months was predicted by negative connectivity between the dACC and the precuneus. Although pain symptoms were positively related to pain symptoms 6 months later, neural predictors were significant after controlling for severity of symptoms at baseline. In effect, results demonstrate that dACC FC is a robust neural predictor of pain outcomes 6 months after injuries associated with MVC.

The findings that positive connectivity between the dACC and the precentral gyrus/primary motor cortex (M1) predicted greater pain severity are consistent with prior literature. For instance, altered stimulation of the M1 in individuals with chronic pain has been widely reported on, related to altered compensatory movements that develop based on injuries.¹⁵ As pain severity at the baseline timepoint was associated with positive




Figure 2. (A) Greater pain severity at baseline is associated with increased functional connectivity between the dACC and precentral gyrus/premotor cortex (peak MNI coordinate: 30, -14, 70). (B) Greater pain severity at 6 months is associated with decreased functional connectivity between the dACC and precuneus (peak MNI coordinate: -6, -74, 32). Display threshold is $P < .001$ whole-brain uncorrected (cluster $P < .05$ false discovery rate-corrected). All effects control for opioid use at baseline, time since injury at baseline, age, and gender. Effects at 6 months additionally control for pain symptoms at baseline and PTSD symptoms at 6 months. dACC, dorsal anterior cingulate cortex; MNI, Montreal Neurological Institute; PTSD, post-traumatic stress disorder.

connectivity between the dACC and M1, this suggests that movement dysfunction, mediated by the M1, may play a greater role in disrupted pain modulation acutely.

We also found that negative FC between the dACC and precuneus predicted greater pain severity 6 months after injury. The precuneus is one of the major “hubs” of the default mode network (DMN), defined as correlated activation between the medial prefrontal cortex, posterior cingulate cortex, and precuneus at rest. Thus, effects specific to the precuneus is unsurprising given the reliance on a resting state paradigm in our design. Increased engagement of the DMN is related to self-referential thinking, planning, and monitoring of internal and external states outside of cognitive or affective demand.¹¹ The precuneus in particular is associated with consolidating information during rest and the forming of cohesive mental representations.¹⁴ Activation of the precuneus diminishes during unconscious states¹⁴ and during the switch from rest to externally focused tasks.³¹

While not traditionally associated with pain processing,²⁸ the precuneus mediates individual differences in pain sensations, or reported conscious experience of pain.³⁶ In healthy individuals, the experience of pain in

the laboratory through the use of controlled thermal heat results in robust engagement of pre-established pain regions, specifically the somatosensory cortex, ACC, and insula.³⁶ However, in these same individuals, subjective report in feeling this pain correlates positively with engagement of the precuneus only (ie, as opposed to other pain-centric regions).³⁶ Other work supports a role for the precuneus in tracking individual differences in pain ratings, but reports opposing effects, such that greater pain sensitivity is related to dis-engagement of the precuneus.⁷⁷ This latter study also demonstrates that greater sensitivity to pain corresponds positively to engagement of the dACC while simultaneously corresponding to disengagement of the precuneus,⁷⁷ suggesting that these brain regions may be part of a functional network. In contrast, there is little evidence that the precuneus is active during pain when perception of pain is not taken into account,³⁶ while there is further evidence that later processing of pain (vs early processing of pain on the order of milliseconds) corresponds to increased precuneus activation.³⁶ Thus, the unfolding of the perception of pain over time as it moves from unconscious to conscious awareness may be moderated by precuneus involvement. The role of the precuneus in information integration and “co-

6 The Journal of Pain

perception," or the integration of our internal and external selves, thus provides the best explanation for the involvement of the precuneus in mediating the internal perception of pain.^{14,46}

This theory with respect to the role of the precuneus in pain processing is further supported by prior research involving those with chronic pain. Patients with chronic pain have increased functional connectivity between the precuneus and thalamus⁴² and precuneus and sensorimotor cortex,⁸ but decreased connectivity between the precuneus and insula.³⁹ Other studies find increased overall DMN connectivity (including the precuneus) with the thalamus in those with chronic pain,⁵¹ suggesting overall strengthening of connections between pain processing and self-referential thinking. Enhanced MPFC-precuneus connectivity and MPFC-thalamic connectivity is directly associated with pain rumination in some of this work,⁵¹ suggesting that precuneus connectivity may be associated with integrating pain into a current state of mind. Finally, other studies have found that patients with chronic pain over-engage the precuneus within the larger DMN, perhaps showing greater reliance on this region during times of rest related to the integration of pain states.⁴ Limited neuroimaging longitudinal studies involving those with chronic pain have been done, although findings from this work also implicate the precuneus. Individuals who experience persistent chronic pain over 1 year exhibit decreased connectivity between the insula and precuneus.⁵ In the present study, variability in the functional connection between the dACC and precuneus was a reliable marker of individual differences in future pain, furthering the role of the precuneus in the perception of pain. Weakened correlation between brain regions that regulate pain (ie, dACC) and those associated with the internal perception of pain (ie, precuneus) may therefore provide a "neuro-profile" of persistent perception of pain.

Contrary to hypotheses, we did not find evidence that dACC-FC with other brain regions involved in pain processing—specifically the brainstem, thalamus, somatosensory cortex, amygdala, and insula—predicted pain severity 6 months after injury. Although these regions are involved in the acute processing of pain, FC between the dACC and these regions was not a useful predictor of chronic pain symptoms. The current analysis controlled for symptoms of PTSD, a disorder associated with affective disturbances involving atypical insula and amygdala engagement.^{30,67} That is, regions involved in

dACC Functional Connectivity and Pain

the affective components of pain perception (eg, insula, amygdala) may also be involved in affective symptoms of PTSD. By controlling for the contribution of stress symptoms in the present analysis, results isolate dACC-FC perturbations specific to the ability to predict chronicity of pain symptoms in comparison to comorbid conditions, which may help explain the lack of findings with other nociceptive regions that have shared disturbances with PTSD.

Results of this study should be considered in light of several limitations. First, a relatively small sample size was used in the present study; therefore, findings require replication. Second, pain was assessed using a singular VAS rating at baseline and 6 months and therefore does not reflect the average pain over a duration of time. Pain severity can be measured both in terms of pain intensity and unpleasantness to further validate pain ratings. Related, this sample experienced relatively lower levels of pain. Future studies should consider incorporating average VAS ratings over a period of time to account for subtle fluctuations in the reporting of pain, include individuals with more severe pain symptoms, and use more robust measures of pain that include ratings of unpleasantness. This investigation is also limited to pain outcomes following MVC and, thus, findings may not be generalizable to other types of injury. As this preliminary investigation focused exclusively on the dACC, future work should also consider investigating other nociceptive regions as predictors of chronic pain. Finally, in this sample, the qualitative nature of the MVC was similar across participants, making it difficult to ascertain whether rsfMRI FC varied as a function of properties of the MVC. Without a control group, we were also unable to conclude that rsfMRI results are directly related to the MVC in this sample.

Despite these limitations, this study provides evidence of distinct neural vulnerabilities that predict chronic pain based on positive versus negative whole-brain connectivity with the dACC. The dACC is implicated in acute pain response^{54,77} and has been viewed as a target for the treatment of chronic pain specifically.⁶⁹ This study provides added evidence of the importance of dACC in pain as a target of treatment by demonstrating that dACC-FC may be able to predict those most at risk for chronic pain. Ultimately, this work supports dACC connectivity as a potential biomarker of chronic pain.

References

1. An X, Bandler R, Ongur D, Price JL: Prefrontal cortical projections to longitudinal columns in the midbrain periaqueductal gray in Macaque monkeys. *J Comp Neurol* 401:455-479, 1998
2. Apkarian AV, Bushnell MC, Treede R-D, Zubieta J-K: Human brain mechanisms of pain perception and regulation in health and disease. *Eur J Pain* 9:463, 2005
3. Asmundson GJ, Coons MJ, Taylor S, Katz J: PTSD and the experience of pain: Research and clinical implications of shared vulnerability and mutual maintenance models. *Can J Psychiatry* 47:930-937, 2002
4. Baliki MN, Mansour AR, Baria AT, Apkarian AV: Functional reorganization of the default mode network across chronic pain conditions. Zang Y-F, editor. *Functional reorganization of the default mode network across chronic pain conditions*. *PLoS One* 9:e106133, 2014
5. Baliki MN, Petre B, Torbey S, Herrmann KM, Huang L, Schnitzer TJ, Fields HL, Apkarian AV: Corticostriatal functional connectivity predicts transition to chronic back pain. *Nat Neurosci* 15:1117-1119, 2012

6. Basbaum AI, Bautista DM, Scherrer G, Julius D: Cellular and molecular mechanisms of pain. *Cell* 139:267-284, 2009
7. Beck JG, Coffey SF: Assessment and treatment of post-traumatic stress disorder after a motor vehicle collision: Empirical findings and clinical observations. *Prof Psychol Res Pr* 38:629-639, 2007
8. Bolwerk A, Seifert F, Maihöfner C: Altered resting-state functional connectivity in complex regional pain syndrome. *J Pain* 14:1107-1115.e8, 2013
9. Bortsov AV, Platts-Mills TF, Peak DA, Jones JS, Swor RA, Domeier RM, Lee DC, Rathlev NK, Hendry PL, Fillingim RB, McLean SA: Pain distribution and predictors of widespread pain in the immediate aftermath of motor vehicle collision: Widespread pain after motor vehicle collision. *Eur J Pain* 17:1243-1251, 2013
10. Brooks J, Tracey I: From nociception to pain perception: Imaging the spinal and supraspinal pathways. *J Anat* 207:19-33, 2005
11. Buckner RL, Andrews-Hanna JR, Schacter DL: The brain's default network: anatomy, function, and relevance to disease. *Ann N Y Acad Sci* 1124:1-38, 2008
12. Bush G, Luu P, Posner MI: Cognitive and emotional influences in anterior cingulate cortex. *Trends Cogn Sci* 4:215-222, 2000
13. Bushnell MC, Čeko M, Low LA: Cognitive and emotional control of pain and its disruption in chronic pain. *Nat Rev Neurosci* 14:502-511, 2013
14. Cavanna AE, Trimble MR: The precuneus: A review of its functional anatomy and behavioural correlates. *Brain* 129:564-583, 2006
15. Chang W-J, O'Connell NE, Beckenkamp PR, Alhassani G, Liston MB, Schabrun SM: Altered primary motor cortex structure, organization, and function in chronic pain: A systematic review and meta-analysis. *J Pain* 19:341-359, 2018
16. Chou R, Shekelle P: Will this patient develop persistent disabling low back pain? *JAMA* 303:1295-1302, 2010
17. Cifre I, Sitges C, Fraiman D, Munoz MA, Balenzuela P, Gonzalez-Roldan A, Martinez-Jauand M, Birbaumer N, Chialvo DR, Montoya P: Disrupted functional connectivity of the pain network in fibromyalgia. *Psychosom Med* 74:55-62, 2012
18. Connolly MR, Chaudari JY, Yang X, Ward N, Kitt RA, Herrmann RS, Krane EJ, LeBel AA, Smith SM, Walco GA, Weisman SJ, Turk DC, Dworkin RH, Gewandter JS: Design and reporting characteristics of clinical trials of select chronic and recurrent pediatric pain conditions: An analgesic, anesthetic, and addiction clinical trial translations, innovations, opportunities, and networks systematic review. *J Pain* 20:394-404, 2019
19. Cook DB, Lange G, Ciccone DS, Liu WC, Steffener J, Natelson BH: Functional imaging of pain in patients with primary fibromyalgia. *J Rheumatol* 31:364-378, 2004
20. Demirakca T, Cardinale V, Dehn S, Ruf M, Ende G: The exercising brain: changes in functional connectivity induced by an integrated multimodal cognitive and whole-body coordination training. *Neural Plast* 2016:1-11, 2016
21. DiMaggio C, Ayoung-Chee P, Shinseki M, Wilson C, Marshall G, Lee DC, Wall S, Maulana S, Leon Pachter H, Frangos S: Traumatic injury in the United States: Inpatient epidemiology 2000–2011. *Injury* 47:1393-1403, 2016
22. Downie WW, Leatham PA, Rhind VM, Wright V, Branco JA, Anderson JA: Studies with pain rating scales. *Ann Rheum Dis* 37:378-381, 1978
23. Duckworth MP, Iezzi T: Chronic pain and posttraumatic stress symptoms in litigating motor vehicle accident victims. *Clin J Pain* 21:251-261, 2005
24. Duenas M, Ojeda B, Salazar A, Mico JA, Failde I: A review of chronic pain impact on patients, their social environment and the health care system. *J Pain Res* 9:457-467, 2016
25. Dum RP, Levinthal DJ, Strick PL: The spinothalamic system targets motor and sensory areas in the cerebral cortex of monkeys. *J Neurosci* 29:14223-14235, 2009
26. Eklund A, Nichols T, Knutsson H: Cluster failure: Why fMRI inferences for spatial extent have inflated false-positive rates. *Proc Natl Acad Sci U S A* 113:7900-7905, 2016
27. Etkin A, Egner T, Kalisch R: Emotional processing in anterior cingulate and medial prefrontal cortex. *Trends Cogn Sci* 15:85-93, 2011
28. Fenton BW, Shih E, Zolton J: The neurobiology of pain perception in normal and persistent pain. *Pain* 22, 2015
29. Ferraz MB, Quresma MR, Aquino LR, Atra E, Tugwell P, Goldsmith CH: Reliability of pain scales in the assessment of literate and illiterate patients with rheumatoid arthritis. *J Rheumatol* 17:1022-1024, 1990
30. Fitzgerald JM, DiGangi JA, Phan KL: Functional neuroanatomy of emotion and its regulation in PTSD. *Harv Rev Psychiatry* 26:116-128, 2018
31. Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME: The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *PNAS* 102:9673-9678, 2005
32. Fuchs PN, Peng YB, Boyette-Davis JA, Uhelski ML: The anterior cingulate cortex and pain processing. *Front Integr Neurosci* 8, 2014. Available at: <http://journal.frontiersin.org/article/10.3389/fnint.2014.00035/abstract>. Accessed August 12, 2020
33. Geuter S, Boll S, Eippert F, Büchel C: Functional dissociation of stimulus intensity encoding and predictive coding of pain in the insula. *eLife* 6, 2017. Available at: <https://elifesciences.org/articles/24770>. Accessed August 12, 2020
34. Ghashghaeia HT, Hilgetag CC, Barbas H: Sequence of information processing for emotions based on the anatomic dialogue between prefrontal cortex and amygdala. *Neuroimage* 34:905-923, 2007
35. Giesecke T, Gracely RH, Grant MAB, Nachemson A, Petzke F, Williams DA, Clauw DJ: Evidence of augmented central pain processing in idiopathic chronic low back pain. *Arthr Rheum* 50:613-623, 2004
36. Goffaux P, Girard-Tremblay L, Marchand S, Daigle K, Whittingstall K: Individual differences in pain sensitivity vary as a function of precuneus reactivity. *Brain Topogr* 27:366-374, 2014
37. Gracely RH: Pain catastrophizing and neural responses to pain among persons with fibromyalgia. *Brain* 127:835-843, 2004

8 The Journal of Pain

38. Herman JP, Ostrander MM, Mueller NK, Figueiredo H: Limbic system mechanisms of stress regulation: Hypothalamo-pituitary-adrenocortical axis. *Prog Neuropsychopharmacol Biol Psychiatry* 29:1201-1213, 2005
39. Hong J-Y, Kilpatrick LA, Labus JS, Gupta A, Katibian D, Ashe-McNalley C, Stains J, Heendeniya N, Smith SR, Tillisch K, Naliboff B, Mayer EA: Sex and disease-related alterations of anterior insula functional connectivity in chronic abdominal pain. *J Neurosci* 34:14252-14259, 2014
40. Hu S, Ide JS, Chao HH, Zhornitsky S, Fischer KA, Wang W, Zhang S, Li CR: Resting state functional connectivity of the amygdala and problem drinking in non-dependent alcohol drinkers. *Drug Alcohol Depend* 185:173-180, 2018
41. Hutchison WD, Davis KD, Lozano AM, Tasker RR, Dostrovsky JO: Pain-related neurons in the human cingulate cortex. *Nat Neurosci* 2:403-405, 1999
42. Ichesco E, Puiu T, Hampson JP, Kairys AE, Clauw DJ, Harte SE, Peltier SJ, Harris RE, Schmidt-Wilcke T: Altered fMRI resting-state connectivity in individuals with fibromyalgia on acute pain stimulation. *Eur J Pain* 20:1079-1089, 2016
43. Iwata K, Kamo H, Ogawa A, Tsuboi Y, Noma N, Mitsuhashi Y, Taira M, Koshikawa N, Kitagawa J: Anterior cingulate cortical neuronal activity during perception of noxious thermal stimuli in monkeys. *J Neurophysiol* 94:1980-1991, 2005
44. Jahn A, Nee DE, Alexander WH, Brown JW: Distinct regions within medial prefrontal cortex process pain and cognition. *J Neurosci* 36:12385-12392, 2016
45. Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM: FSL. *Neuroimage* 62:782-790, 2012
46. Johnson MK, Raye CL, Mitchell KJ, Touryan SR, Greene EJ, Nolen-Hoeksema S: Dissociating medial frontal and posterior cingulate activity during self-reflection. *Soc Cogn Affect Neurosci* 1:56-64, 2006
47. Jones GT, Nicholl BI, McBeth J, Davies KA, Morriss RK, Dickens C, Macfarlane GJ: Role of road traffic accidents and other traumatic events in the onset of chronic widespread pain: Results from a population-based prospective study. *Arthr Care Res* 63:696-701, 2011
48. Kirino E, Tanaka S, Fukuta M, Inami R, Arai H, Inoue R, Aoki S: Simultaneous resting-state functional MRI and electroencephalography recordings of functional connectivity in patients with schizophrenia: rsfMRI-EEG of FC in schizophrenia. *Psychiatry Clin Neurosci* 71:262-270, 2017
49. Kong J, Tu P, Zyloney C, Su T: Intrinsic functional connectivity of the periaqueductal gray, a resting fMRI study. *Behav Brain Res* 211:215-219, 2010
50. Koyama T, Tanaka YZ, Mikami A: Nociceptive neurons in the macaque anterior cingulate activate during anticipation of pain. *Neuroreport* 9:2663-2667, 1998
51. Kucyi A, Moayed M, Weissman-Fogel I, Goldberg MB, Freeman BV, Tenenbaum HC, Davis KD: Enhanced medial prefrontal-default mode network functional connectivity in chronic pain and its association with pain rumination. *J Neurosci* 34:3969-3975, 2014
52. Legrain V, Iannetti GD, Plaghki L, Mouraux A: The pain matrix reloaded. *Prog Neurobiol* 93:111-124, 2011
53. Liddell BJ, Brown KJ, Kemp AH, Barton MJ, Das P, Peduto A, Gordon E, Williams LM: A direct brainstem-amygdala-cortical 'alarm' system for subliminal signals of fear. *Neuroimage* 24:235-243, 2005
54. Lieberman MD, Eisenberger NI: The dorsal anterior cingulate cortex is selective for pain: Results from large-scale reverse inference. *Proc Natl Acad Sci* 112:15250-15255, 2015
55. Maldjian JA, Laurienti PJ, Kraft RA, Burdette JH: An automated method for neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI data sets. *Neuroimage* 19:1233-1239, 2003
56. McCormack HM, Horne DJ, Sheather S: Clinical applications of visual analogue scales: A critical review. *Psychol Med* 18:1007-1019, 1988
57. McLean SA, Ulirsch JC, Slade GD, Soward AC, Swor RA, Peak DA, Jones JS, Rathlev NK, Lee DC, Domeier RM, Hendry PL, Bortsov AV, Bair E: Incidence and predictors of neck and widespread pain after motor vehicle collision among US litigants and nonlitigants. *Pain* 155:309-321, 2014
58. Menon V, Uddin LQ: Saliency, switching, attention and control: a network model of insula function. *Brain Struct Funct* 214:655-667, 2010
59. Mertens P, Blond S, David R, Rigoard P: Anatomy, physiology and neurobiology of the nociception: A focus on low back pain (part A). *Neurochirurgie* 61:S22-S34, 2015
60. Morasco BJ, Lovejoy TI, Lu M, Turk DC, Lewis L, Dobscha SK: The relationship between PTSD and chronic pain: Mediating role of coping strategies and depression. *Pain* 154:609-616, 2013
61. Morecraft RJ, McNeal DW, Stilwell-Morecraft KS, Gedney M, Ge J, Schroeder CM, van Hoesen GW: Amygdala interconnections with the cingulate motor cortex in the rhesus monkey. *J Comp Neurol* 500:134-165, 2007
62. Peyron R, Laurent B, Garcia-Larrea L: Functional imaging of brain responses to pain. A review and meta-analysis. *Clin Neurophysiol* 30:263-288, 2000
63. Poldrack RA, Baker CI, Durnez J, Gorgolewski KJ, Matthews PM, Munafò MR, Nichols TE, Poline J-B, Vul E, Yarkoni T: Scanning the horizon: Towards transparent and reproducible neuroimaging research. *Nat Rev Neurosci* 18:115-126, 2017
64. Power JD, Mitra A, Laumann TO, Snyder AZ, Schlaggar BL, Petersen SE: Methods to detect, characterize, and remove motion artifact in resting state fMRI. *Neuroimage* 84:320-341, 2014
65. Power JD, Schlaggar BL, Petersen SE: Recent progress and outstanding issues in motion correction in resting state fMRI. *Neuroimage* 105:536-551, 2015
66. Pujol J, López-Solà M, Ortiz H, Vilanova JC, Harrison BJ, Yücel M, Soriano-Mas C, Cardoner N, Deus J: Mapping brain response to pain in fibromyalgia patients using temporal analysis of fMRI. García AV, editor. Mapping brain response to pain in fibromyalgia patients using temporal analysis of fMRI. *PLoS One* 4:e5224, 2009
67. Rauch SL, Shin LM, Phelps EA: Neurocircuitry models of posttraumatic stress disorder and extinction: Human neuroimaging research—Past, present, and future. *Biol Psychiatry* 60:376-382, 2006

68. Roy AK, Shehzad Z, Margulies DS, Kelly AMC, Uddin LQ, Gotimer K, Biswal BB, Castellanos FX, Milham MP: Functional connectivity of the human amygdala using resting state fMRI. *Neuroimage* 45:614-626, 2009
69. Russo JF, Sheth SA: Deep brain stimulation of the dorsal anterior cingulate cortex for the treatment of chronic neuropathic pain. *Neurosurg Focus* 38:E11, 2015
70. Salomons TV, Johnstone T, Backonja M-M, Davidson RJ: Perceived controllability modulates the neural response to pain. *J Neurosci* 24:7199-8203, 2004
71. Schmidt-Wilcke T, Ichesco E, Hampson JP, Kairys A, Peltier S, Harte S, Clauw DJ, Harris RE: Resting state connectivity correlates with drug and placebo response in fibromyalgia patients. *Neuroimage* 6:252-261, 2014
72. Shackman AJ, Salomons TV, Slagter HA, Fox AS, Winter JJ, Davidson RJ: The integration of negative affect, pain and cognitive control in the cingulate cortex. *Nat Rev Neurosci* 12:154-167, 2011
73. Thompson JM, Neugebauer V: Amygdala plasticity and pain. *Pain Res Manag* 2017:1-12, 2017
74. Tracey I, Mantyh PW: The cerebral signature for pain perception and its modulation. *Neuron* 55:377-391, 2007
75. Trevino C, Harl F, deRoon-Cassini T, Brasel K, Litwack K: Predictors of chronic pain in traumatically injured hospitalized adult patients. *J Trauma Nurs* 21:50-56, 2014
76. Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N, Mazoyer B, Joliot M: Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* 15:273-289, 2002
77. Wager TD, Atlas LY, Lindquist MA, Roy M, Woo C-W, Kross E: An fMRI-Based neurologic signature of physical pain. *N Engl J Med* 368:1388-1397, 2013
78. Weiss DS, Marmar CR: The impact of events scale—Revised. *Assessing Psychological Trauma and PTSD*. New York, NY, Guilford, 1996, pp 399–411
79. Whitfield-Gabrieli S, Nieto-Castanon A: Conn: A functional connectivity toolbox for correlated and anti-correlated brain networks. *Brain Connect* 2:125-141, 2012
80. Woo C-W, Krishnan A, Wager TD: Cluster-extent based thresholding in fMRI analyses: Pitfalls and recommendations. *Neuroimage* 91:412-419, 2014
81. Yu C, Zhou Y, Liu Y, Jiang T, Dong H, Zhang Y, Walter M: Functional segregation of the human cingulate cortex is confirmed by functional connectivity based neuroanatomical parcellation. *Neuroimage* 54:2571-2581, 2011